

Laboratory Preparation of the Components of a Dye-Sensitized Solar Cell

Ezinna Lucky Efurumibe¹, Anyalewechi Daniel Asiegbu² and Michael Onuu³

1. Physics Department, College of Natural and Physical Sciences, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria.
2. Physics Department, College of Natural and Physical Sciences, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria.
3. Physics Department, University of Calabar, Calabar
efurumibeezinna@yahoo.com

Abstract

In this laboratory preparation of the components, the solutions of some of the components of the cell were prepared. This was done (in some cases) after measuring out certain amount of the chemicals. For example for the TiO_2 , 6grams of the powder was first weighed, after which the solution was prepared. At the end of the laboratory session, a solution (suspension) of TiO_2 was prepared. Afterwards, a solution of anthocynin dye from *Momordica Charantia* seed was obtained separately.

1. INTRODUCTION

Energy is inevitably the most wanted commodity in the world today. The industrial revolution has increased the quest for energy, particularly energy whose means of production is environmental friendly. Dye-sensitized solar cell is an interesting area of study for many scientists because of its promising features (O'Regan and Grätzel, 1991). Number one feature is that it only needs a little light from the sun to be activated. Number two feature is that its means of energy production is environmentally friendly. It does not require so much capital and technology to fabricate. Before a dye-sensitized solar cell is built/coupled, its components need to be prepared first. Such components include: the anode of the cell, the sensitized dye and the electrolyte of the cell (Abayev *et al*, 2003; Grätzel, 2004). In most cases the preparation requires first of all getting the solution of the components; and by some other processes, the components of the cell can be built. Here we present majorly the laboratory preparation of TiO_2 suspension and anthocynin solution from *Momordica Charantia* seed. These solutions were prepared in the Chemistry and Biochemistry laboratories of Michael Okpara University of Agriculture, Umudike. The aim of the exercise is to produce a suspension of TiO_2 and a solution of anthocyanin dye.

2. PREPARING THE TiO_2 SUSPENSION

In preparing the TiO_2 suspension, 9ml of nitric acid solution (pH 3.04 in deionized water) was added (at 1ml increment) to 6g of TiO_2 powder in a mortar and grounded (with a pestle) until the powder dissolved in the solution (Fanis, 2010). The grinding was done for about 30 minute after which 3 drops of liquid detergent was added to the solution to make for even coating of the TiO_2 suspension on the conductive glass slides (see fig. 1). After which the solution was stored in a dropper bottle for coating the conductive glass slides.



Fig.1: A digital balance used in measuring 6g of TiO_2 powder

3. PREPARING THE ANTHOCYANIN FROM MOMORDICA CHARANTIA SEED

The organic dye used in staining the coated glass slides was ‘anthocyanin’ obtained from the seed of *Momordica Charantia* (see fig. 2). The dye was obtained by crushing the redish-yellow seed of the plant (*Momordica Charantia*) in a mortar with the addition of acetone (see fig. 3). The acetone helped in producing the solution of the dye (Greenwood and Earnshaw, 1984).



Fig.2: A lab scientist preparing the deionized water for the TiO_2 suspension



Fig.3: Preparing the TiO_2 suspension

4. RESULT

At the end of the laboratory exercise, TiO_2 suspension was obtained. This was placed inside a dropper bottle for later use. This suspension was later used to coat some glass slides for the purpose of designing a dye-sensitized solar cell. The next component that was obtained from the laboratory exercise was an organic dye, which was the anthocynin from the Momordica Charantia seed. This anthocynin was used to stain the glass slides (see fig.4), in the course of designing dye-sensitized solar cells.



Fig. 4: Seed of Momordica Charantia



Fig. 5: Grinding the Momordica Charantia seed

5. CONCLUSION

Working with chemicals in the laboratory is a wonderful experience, particularly when the aim is to build a physical device. Dye sensitized solar cell, as was said in one of our papers is a new research area; an area for prospective researcher in the field of Physics and Engineering (Efurumibe *et al*, 2012). It our belief that one day a good organic dye will be discovered which would be capable of boosting the efficiency of dye-sensitized solar cell up to 80% at least. Nevertheless for the study at hand, the aim of the laboratory exercise was achieved: A suspension of TiO_2 was obtained. Also was obtained is a solution of anthocynin dye, for design of a dye-sensitized solar cell.

6. ACKNOWLEDGEMENT

I would not forget the following persons: Uchenna Ariwa (of Biochemistry Department), Kanu I. Kalu (of Chemistry Department), Solomon Nwadinobi (of Chemistry Department) and Ngozi Ukanwaolu (Physics Department) for their contributions towards the realization of the aim of this work. They helped in providing the deionized water, nitric acid solution and relevant apparatuses for the work.

REFERENCES

- Abayev, I., Zaban, A., Fabregat-Santiago, F and Bisquert, J. (2003). Electronic Conductivity in nanostructured TiO_2 films permeated with electrolyte, *Physica Status Solidi (a)* 196, pp 4-6
- Efurumibe, E.L., Asiegbu, A.D. and Onuu, M.U. (2012). Mathematical Modelling of Electron Transport through the Anode (TiO_2) of a standard dye-sensitized solar cell, *Asian Journal of Applied Sciences*, pp 33-44

Fanis, L. (2010). Nanocrystalline solar cell kit Recreating Photosynthesis, ICE publication, pp52

Grätzel, M. (2004). Conversion of sunlight to electric power by nanocrystalline dye-sensitized solar cells, Journal of photochemistry and photobiology A: Chemistry, 164 (1): 3-14

Greenwood, N., Earnshaw, A. (1984), Chemistry of the Elements, Oxford: Pergamon, pp. 1117–1119, accessed online on 16/07/2011 from: <http://webcache.googleusercontent.com>

O'Regan, B. and Grätzel, M. (1991). A low cost, high-efficiency solar cell based on dye-sensitized colloidal TiO₂ films, Nature 353, pp 737-739

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

CALL FOR PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request from readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

